

LOCALIZED AMYLOIDOSIS OF LARYNX: A CASE REPORT AND REVIEW OF RECENT CASES

LARİNKSİN LOKALİZE AMİLOİDOZU: VAKA RAPORU VE SON YAYINLANAN VAKALARIN DEĞERLENDİRİLMESİ Laringoloji Başvuru: 20.03.2025 Kabul: 23.06.2025 Yayın: 23.06.2025

Ali Ozan Yazıcı¹, Güler Berkiten², Belgin Tutar³, Yavuz Uyar²

¹ Sağlık Bilimleri Üniversitesi Prof. Dr. Cemil Taşcıoğlu Şehir Hastanesi Kulak Burun Boğaz Kliniği ² Okmeydanı Eğitim ve Araştırma Hastanesi ³ Okmeydanı Eğitim ve Araştırma Hstanesi

Özet

Amiloidoz vücudun çeşitli organlarında amiloid proteinin birikimiyle karakterize bir hastalıktır. Sistemik görülmesi lokalize görülmesine göre fazla sıklıkta olmasıyla birlikte larinksin lokalize tutulumu oldukça nadir görülmektedir. Bu olguda kliniğimize ses kısıklığı ile başvuran 59 yaşında erkek hasta ele alınmıştır. Laringoskopik muayenesinde vokal kordların anterior komüssüründe, epiglotun petiolünde ve subglottik alanda düzensiz kitleler görülen hastadan çoklu biyopsiler alınmıştır. Histopatolojik sonuç, kongo red ve kristal viyole pozitifliği amiloid ile uyumlu gelmiştir. Her ne kadar lokal amiloidoz nadir görülse de larinks kitlelerinde ayırıcı tanıda düşünülmesi gerekmektedir.

Anahtar kelimeler: Disfoni, Amiloidoz Larinks

Abstract

Amiloidosis is a diasese characterized by accumilation of amyloid protein on several organs of the body. Although systemic involvement is more common, localized laryngeal involvement quite rare. In this case, a 59-year-old male patient presenting with hoarseness to our clinic was discussed. After laryngoscopic examination, multiple biopsies were taken from irregular were observed masses on anterior commissure of vocal cord, the petiole of epyglottis and subglottic area. Histopatological findings were consistent with amyloid deposition, incluiding Congo Red and crystal violet positivity. Whilst localized amyloidosis is rare, it should be considered as a differential diagnosis of larygngeal masses.

Keywords: Dysphonia, Amyloidosis Larynx

Introduction

Amyloidosis is a group of diseases characterized by the deposition of abnormal proteins (insoluble fibrillar proteins) in the extracellular space of the body. The accumulation can be systemic or localized [1]. While amyloid accumulation is possible in several organs of the body (heart, liver, kidney, spleen, etc.), laryngeal involvement is exceedingly rare [2]. Localized laryngeal amyloidosis accounts for 0.2–1.2% of all benign laryngeal tumors. Nevertheless, the larynx remains one of the most commonly affected organs in the neck region. It is more common in males than in females and peaks in the fourth to sixth decades of life [3]. Although symptoms of respiratory amyloidosis are nonspecific, they may include cough, dyspnea, wheezing, hemoptysis, and dysphonia [4]. In this case, a patient presenting with dysphonia was evaluated. Furthermore, the data extracted from the reviewed articles including age, gender, symptoms, laryngeal localization of amyloidosis, treatment strategies, and follow-up durations are discussed in conjunction with the present case and summarized in Table 1.

Sorumlu Yazar: Ali Ozan Yazıcı, Sağlık Bilimleri Üniversitesi Prof. Dr. Cemil Taşcıoğlu Şehir Hastanesi Kulak Burun Boğaz Kliniği Kaptanpaşa Mahallesi Darülaceze Cad. No:27 Şişli aliiozany@gmail.com



	Age	Gender	Symptom	Localization	Treatment	Follow-up Periods and Recurrence
Garefis et al, 2023 (9)	73	Male	Dysphonia	Right vocal cord and left vestibular fold	Laser CO2	2 years (no recurrence)
Feng et al, 2022 (10)	51	Male	Dysphonia	Left vocal cord and nasopharyux	Plasma knife	3th and 6th months (no recurrence)
Choudhary and Koldradkar, 2021 (1)	22	Male	Dysphonia	Left false cord, ipsilateral arytenoid and ventricle	Coblator Laryngeal Wand	6 months (no recurrence)
Zhao and Liu, 2021 (11)	43	Female		Right vocal cord, subglottic area	Cold Technique	2nd month (no recurrence)
Ghauth et al, 2021(12)	50	Male	Dysphonia	Right false cord	Cold Technique	1st month (with recurrence)
Mesolella et al, 2020 (13)	14	Female	Dysphonia	Left false cord	Laser CO2	Monthly in first 6 months, bimonthly until the end of second year, six-monthly for 2 years, yearly from the fifth year) (no recurrence)
	21	Female	Dysphonia, vocal fatigue	Right false cord and aryepiglottic fold	Laser CO2	Recurrence on 1st month, after second operation 5th year with no recurrence
	34	Female	Dysphonia, stridor	Subglottic	Laser CO2	Monthly in first 6 months, bimonthly until the end of second year, six-monthly for 2 years, yearly from the fifth year) (no recurrence)
	47	Female	Dysphonia, dyspnea	Diffuse laryngeal involvement	Laser CO2	Operated 3 times, no recurrence
	38	Female	Dysphonia	Left false cord	Cold Technique	Operated 2 times, 1st year with no recurrence
Stavrakas et al, 2019 (14)	27	Male	Dysphonia	Left aryepiglottic fold, left false fold, right arytomoid area	Cold Technique	6 monthly follow-up, no recurrence after 2

 Table 1

 A Five-Year Review of Recent Cases

Case Report

A 59 year old male patient presented to our clinic with a complaint of hoarseness that had persisted for 4 months, without any additional symptoms such as dyspnea, cough, or dysphagia. The patient had no known comorbidities or family history and had a history of 30 pack-years of smoking. On physical examination, otoscopy, rhinoscopy, and intraoral examination appeared normal, while laryngoscopic examination revealed irregular masses on the anterior commissure of the vocal cords, the petiole of the epiglottis, and the subglottic area (Figure 1), which led to the decision for a biopsy.



Figure 1

Laryngoscopic examination of the patient, irregular masses on the anterior commissure of the vocal cords, the petiole of the epiglottis, and the subglottic area.

Histopathological analysis of the multiple biopsies revealed amorphous material compatible with amyloid in the subepithelial area, confirmed by positive Congo red and crystal violet staining (Figure 2,3).

ENTCase



Figure 2 Pathological images of the collected samples. Pink amorphous material in subepithelial area stained by Haematoxylin & Eosin (at 20x magnification).



Figure 3A Amyloid stain with Congo red (A, B). A, material consistent with subepithelial amyloid (at 20x magnification).

ENTCase



Figure 3B

Amyloid stain with Congo red (A, B). B, accumulation of material consistent with amyloid in vessel wall (at 100x magnification).



Figure 3C C, apple green birefringence under polarized light microscopy.

To differentiate between systemic and localized amyloidosis, further investigations were requested through an internal medicine consultation.

The patient's laboratory results, including complete blood count, liver and kidney function tests, serum calcium, potassium, erythrocyte sedimentation rate, and serum immunoglobulins, were within normal limits. Protein levels in serum and urine were assessed, and both were found to be normal. No monoclonal proteins were detected in serum and urine protein electrophoresis. The electrocardiogram and chest X-ray were normal. After excluding diseases that could cause systemic amyloidosis, it was determined that the amyloidosis was localized to the larynx. Given the improvement in the patient's symptoms, close follow-up and continued monitoring were deemed appropriate for the treatment plan. Follow-up evaluations were performed at the first, third, and sixth months, during which endoscopic examinations demonstrated no signs of lesion progression or the emergence of new formations.

ENTCase

Discussion

Amyloidosis is a disease characterized by the accumulation of insoluble abnormal proteins in the extracellular space, resulting in dysfunction across various organs. This condition, which may be either systemic or localized, is relatively rare in the larynx [5]. Although its etiology is not fully understood, several theories propose that plasma cells may produce amyloid light chains, or that there may be deficiencies in the degradation of these light chains within the body [6]. In certain cases, genetic mutations have also been identified as a contributing factor [7].

Depending on whether the condition is systemic or localized, the type of amyloid protein involved may vary. The most commonly encountered forms include AL (amyloid derived from immunoglobulin light chains; the most frequent type), AA (amyloid derived from serum amyloid A protein), ATTR (amyloid derived from the transport protein transthyretin), and dialysis-related amyloidosis (amyloid composed of beta-2 microglobulin) [8].

In systemic amyloidosis, amyloid deposits accumulate in organs such as the liver, heart, spleen, and kidneys, leading to clinical manifestations including hepatomegaly, cardiomegaly, proteinuria, and macroglossia [8]. Localized laryngeal amyloidosis accounts for 0.2–1.2% of all benign laryngeal tumors. However, the larynx is the most commonly affected organ in the head and neck region in cases of amyloidosis [1].

A review of eleven laryngeal amyloidosis cases published in the last five years showed that most of the patients had amyloid deposits localized to the glottis (Table 1). In our case, deposits were observed not only in the glottis but also in the supraglottic and subglottic regions. The mean age of the patients was 38 years, ranging from 14 to 73 years. The study included five male and six female patients [1,9–13].

In laryngeal amyloidosis, the most common symptom is hoarseness, although some patients may present with dyspnea, stridor, persistent dry cough, sleep apnea syndrome, or dysphagia [1]. A review of recent cases demonstrated that the primary symptom of laryngeal amyloidosis was dysphonia, with stridor and dyspnea also reported as accompanying findings [1,9–13,14]. Our patient likewise presented to our clinic with dysphonia.

On laryngoscopic examination, amyloid deposition may appear in various forms, including nodular, polypoid, or diffuse infiltration. The differential diagnosis should include neoplastic diseases, fibroma, papilloma, and tuberculosis.

The gold standard for diagnosis is histopathological analysis, in which amyloid deposits are stained with Congo red and exhibit apple-green birefringence under polarized light microscopy [5]. Following the diagnosis, further investigations including complete blood count, serum calcium, sodium, and potassium levels, liver and kidney function tests, serum immunoglobulin levels, serum and urine protein electrophoresis, electrocardiogram, and chest X-ray should be performed to determine whether the condition is systemic or localized.

Localized amyloidosis is generally considered benign, and its management is guided by the patient's symptoms. The main goal is to maintain adequate airway patency and preserve the functional larynx. Treatment may vary from simple observation, more conservative endoscopic excision to radical excision by partial laryngectomy depending upon the extent of involvement of larynx. Endoscopic excision can be done either with cold knife, CO2 Laser, Microdebrider or with Coblation wands [1]. In cases with large obstructive lesions or anticipated postoperative laryngeal edema, tracheostomy may be required. Small, asymptomatic lesions should be monitored closely [15]. The review indicates that the most commonly employed treatment method was CO_2 laser, accounting for 45% of the eleven reported cases.

A well-defined follow-up process is essential for effective management. The overall prognosis of Laryngeal Amyloidosis is generally good, but the probability of recurrent or residual disease is significant [16]. In the event



of recurrence, repeated resections may be necessary [15].

In published case series involving ten or more patients with clearly defined localized laryngeal amyloidosis and a follow-up period exceeding five years, none required revision surgery beyond the seventh year. Therefore, it is advisable to maintain laryngological follow-up for at least seven years after the last surgical procedure in order to monitor for potential recurrences [16]. Mesolella M, and al. recommend monthly endoscopic check-ups for the first six months, then bimonthly until the end of the second year, then six-monthly for two years and yearly from the fifth year [13]. In the five-year review of recent cases, the mean follow-up duration was reported as 33 months (2.75 years), with two patients exhibiting recurrence during the first-month follow-up [1,9–14]. In our case, due to the presence of small lesions and improvement in the patient's symptoms, a strategy of close follow-up and continuous monitoring was deemed appropriate for the treatment plan. Follow-up assessments were conducted at the first, third, and sixth months, and endoscopic examinations revealed no evidence of mass progression or development of new lesions.

References

- Choudhary SV, Kolwadkar N. Primary Localized Amyloidosis of Larynx: A Case Report. Indian J Otolaryngol Head Neck Surg. 2022;74(Suppl 3):5000-5002. doi:10.1007/s12070-021-02584-y
- 2. Thuong Vu L, et al. Laryngotracheobronchial amyloidosis: A case report. Respir Med Case Rep. 2021;32:101377. Published 2021 Mar 4. doi:10.1016/j.rmcr.2021.101377
- 3. Gallivan GJ, Gallivan HK. Laryngeal amyloidosis causing hoarseness and airway obstruction. J Voice. 2010;24(2):235-239. doi:10.1016/j.jvoice.2008.07.006
- 4. Uyaguari JP, Quizhpe PJ, Lima E. Laryngo-tracheobronchial amyloidosis. Braz J Otorhinolaryngol. 2020;86(6):825-826. doi:10.1016/j.bjorl.2020.07.006
- 5. Kapoor M, et al. Clinical Presentation, Diagnosis and Treatment of TTR Amyloidosis. J Neuromuscul Dis. 2019;6(2):189-199. doi:10.3233/JND-180371
- 6. Berg AM, et al. Localized amyloidosis of the larynx: evidence for light chain composition. Ann Otol Rhinol Laryngol. 1993;102(11):884-889. doi:10.1177/000348949310201112
- Ihne S, et al. Amyloidosis-the Diagnosis and Treatment of an Underdiagnosed Disease. Dtsch Arztebl Int. 2020;117(10):159-166. doi:10.3238/arztebl.2020.0159
- 8. Bustamante JG, Zaidi SRH. Amyloidosis. In: StatPearls. Treasure Island (FL): StatPearls Publishing; July 31, 2023.
- 9. Garefis K, et al. Laryngeal Amyloidosis: Occurrence Long After Radiation Therapy for Laryngeal Cancer. Ear Nose Throat J. Published online June 2, 2023. doi:10.1177/01455613231178117
- 10. Feng Y, et al. Amyloidosis in the nasopharynx and larynx: a case report. J Int Med Res. 2022;50(12):3000605221144151. doi:10.1177/03000605221144151
- 11. Zhao H, Liu C. Laryngeal amyloidosis. Can J Anaesth. 2022;69(4):550-551. doi:10.1007/s12630-021-02179-9
- 12. Ghauth S, et al. Laryngeal amyloidosis: a rare etiology of hoarseness. QJM. 2022;114(12):889-890. doi:10.1093/qjmed/hcab256
- 13. Mesolella M, et al. Focus on localized laryngeal amyloidosis: management of five cases. Open Med (Wars). 2020;15(1):327-332. Published 2020 Apr 20. doi:10.1515/med-2020-0400
- 14. Stavrakas M, et al. Laryngeal Amyloidosis: An Interesting Presentation of Voice Quality Deterioration. Ear Nose Throat J. 2020;99(2):128-129. doi:10.1177/0145561319837473
- 15. Mira C, et al. Localised laryngotracheal amyloidosis: a differential diagnosis not to forget. BMJ Case Rep. 2021;14(2):e237954. Published 2021 Feb 1. doi:10.1136/bcr-2020-237954
- 16. Hazenberg AJ, Hazenberg BP, Dikkers FG. Long-term follow-up after surgery in localized laryngeal



amyloidosis. Eur Arch Otorhinolaryngol. 2016;273(9):2613-2620. doi:10.1007/s00405-016-4061-y